

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 437 334 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
14.07.2004 Bulletin 2004/29

(51) Int Cl.7: **C05F 11/10**

(21) Application number: **04290051.4**

(22) Date of filing: **08.01.2004**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**
Designated Extension States:
AL LT LV MK

(30) Priority: **08.01.2003 CL 242003**

(71) Applicant: **Universidad de los Lagos
Osorno (CL)**

(72) Inventors:
• **De La Fuente Jimenez, Lucia Elena
Osorno (CL)**
• **Santibanez Handschuh, Alejandro Eugenio
Puerto Varas (CL)**

(74) Representative: **Hubert, Philippe et al
Cabinet Beau de Loménie
158, rue de l'Université
75340 Paris Cédex 07 (FR)**

(54) **Productive process for manufacturing an algal species-based organic complement for vegetal fertilization**

(57) An organic complement for fertilizing vegetable species, as an organic product which fulfills with the features of a growth stimulator, mainly manufactured from green algae (*Ulva rigida*) and brownish algae (*Macrocystis pyrifera*). The invention includes a process for obtaining the complement that requires washing, grinding, acid and alkaline digestion, filtering and packaging steps. The most important used supplies are hydrochlo-

ric and phosphoric acid, and potassium carbonate. The product allows to improve the absorption efficiency of the nutrients supplied by the fertilizers, guaranteeing an optimal growth for the plant species, achieving a higher production in a short period of time. The product is biodegradable and beneficial for soils.

EP 1 437 334 A1

Description**Cross-Reference to Related Applications**

- 5 [0001] The instant application claims priority over the Chilean Patent application CL 24-2003, filed on January 8, 2003.

Background of the invention

- 10 [0002] The increased demand for organic fertilizers in developed countries and the current tendency to its sustainability regarding the industrial wastes management, which minimizes the environmental impacts, are situations considered as relevant. In this context, it is envisaged to provoke a positive environmental impact generating a productive alternative directed to the use of low commercial value algae for the production of a complement for organic fertilization.

Brief summary of the invention

- [0003] The features of this organic growth complement are based on function of the marine algae contributions, which comprise a wide range of mineral elements, aminoacids, vitamins and auxin- and cytokinin-type phytohormones.
- 20 [0004] The resulting product contains these elements in an active form; therefore, a higher plant production and/or better yielding can be obtained when it is supplemented to plant crops with a basal fertilization suitable for the vegetal type. Furthermore, this product fulfills meets the following organic standards: EU N° 2092/1; USDA/NOP - Final rule (USA), and the JAS Japanese Agricultural standard for Organic Agricultural Products (Japan).
- [0005] This organic complement for basal fertilization allows better seed germination, an increase in root development, a faster and more uniform plant stabilisation, an increase in the nutrient absorption, a more efficient nutrients
- 25 exploitation, tissue composition improvements, higher resistance to frosts, higher resistance to drought and a faster recovery, higher resistance to diseases and plagues (by fungi and insects), and longer shelf-life.
- [0006] The term "basal fertilization" must be understood as the soil fertilization necessary to be applied to a soil once checked the soil needing of macronutrients (nitrogen, phosphorus and potassium) and micronutrients (molybdenum, magnesium, boron, etc.) considering the vegetal type to be growth.
- 30 [0007] When applying this product as a fertilization complement achieves a maximal effect within a monitored fertilization program, with a constant soil analysis. Using this product, soils do not require (as usually), an over-dosage, since a minimal but exact fertilizer contribution is enough in order to achieve optimal yields.
- [0008] Furthermore, it is important to note that the obtained "organic product" as prepared from raw organic matter consisting of "organic" sources, through the process herein described, allows to provide a fertilizer complement for
- 35 plant growing certifiable for being used for organically grown food crops.

BRIEF DESCRIPTION OF THE DRAWINGS

- [0009] **Figure 1:** shows a flow chart for a typical process for making the organic fertilizer complement.
- 40 [0010] **Figure 2:** shows a mass balance and process condition flow chart for the manufacture of an organic fertilizer complement embodiment of the invention.
- [0011] **Figure 3:** shows a typical processing equipment system for producing the organic complement.

DETAILED DESCRIPTION OF THE INVENTION

- 45 [0012] The present invention provides an organic fertilizer complement, its method of making the fertilizer complement. The organic fertilizer complement is manufactured from green algae (*Ulva rigida*, hereinafter "*ulva*") and brownish algae (*Macrocystis pyrifera*, hereinafter "*macrocystis*"). The organic product provides features of a vegetal growth stimulator for plants. The product provides improved absorption efficiency of the nutrients supplied by the fertilizers,
- 50 guaranteeing an optimal growth for the plant species, and achieving higher production in a short period of time. the product is biodegradable and beneficial for soils.
- [0013] The process for obtaining the fertilizer complement from the *ulva* and *macrocystis* typically includes the steps of washing, grinding, acid and alkaline digestion, filtering and packaging. typical other raw materials are inorganic acids and salts, such as hydrochloric and phosphoric acid, and potassium carbonate.
- 55

- **Raw Materials Reception:** Fresh, whole *Macrocystis*, previously washed with salted water in order to remove sand and foreign organisms, is received. *Ulva* is received, dried and minced. Both raw materials must be in good conditions and without malodors or other strange odors. Any equipment items used in the description below are

shown in parenthesis and referenced in Figure 3.

- **Grinding:** The dried *Ulva* is ground to about 0.2 mm in a blade mill (12) for achieving good homogenization when later admixed with the *Macrocystis* during the alkaline digestion step, described below.


- **Weighing:** The fresh alga *Macrocystis* and the ground and dried alga *Ulva* are weight into portions at a weight ratio of 1:1 on a fresh algae base. The weight ratio can vary up to 10:1 if dried *Ulva* is used.

- **Washing:** The fresh alga *Macrocystis* is placed in the perforated basket (16), and immersed into a conical vessel (1) containing cold tap water. This operation is repeated until it is established that the *Macrocystis* is clean and that the salts have been removed satisfactorily. A checking of the salt removal can typically be carried out by pouring about 20 ml of washing liquid in a tube and adding 3 drops of AgNO_3 . If a white precipitate forms, then excessive salts are still present. This operation and the checking are carried out until the precipitate formed is minimal. Alternative means of checking can include a batch or continuous analysis of the electric conductivity of the washing liquid using instrumentation that are well known to those skilled in the art. For accelerating the washing process at the time of draining the vessel, water can be sprayed.

- **Mincing:** The washed alga *Macrocystis* is then minced in a turbine mill (13) until a typical particle size of about 1 cm diameter is achieved. The minced algae is subsequently placed in a perforated basket (16). This step has as a first objective to facilitate the handling of the alga during the process, and as a second objective to achieve a particle size which allows the subsequent reagents to uniformly penetrate within the alga, thus achieving more homogeneous reactions and better control of the chemical treatment.

- **Acid Treatment:** The perforated basket with the minced *Macrocystis* is immersed in the jacketed vessel (2) containing, per 1 Kg of *Macrocystis*, 2 parts by weight of water and 25 ml of 0.2 N HCl, for 30-40 minutes at $\pm 50^\circ\text{C}$ under constant stirring (mixer 17). This treatment removes the mineral salts and the excess soluble organic material which was not removed in the previous washing step, as well as the sediments and organisms associated with the raw algae. An ionic exchange chemical reaction is carried out, principally between the calcium ions and other divalent ions as magnesium and strontium contained in the algae as the corresponding divalent metal alginate, thus producing alginic acid (HAlg).

[0014] The chemical reaction is as follows:




- **Draining:** To carry out the residual liquid and solid separation, the perforated basket (16) is raised out of the jacketed vessel (2), leaving the liquid to drain by gravity.

- **Washing:** The resulting drained solids are washed with cold water, such as by spraying water through a hose into the basket containing the algae, for complete removal of the alginic acid.

- **Grinding:** The washed algae *Macrocystis* is ground in the turbine mill (13) having a sieve that has 0.4 cm holes, for mechanically reducing the algal particles to assist in their dissolution during the subsequent digestion step; thus, the smaller the particle size, the more effective shall be the reaction.

- **Digestion:** An alkalinizing agent, potassium carbonate salt (K_2CO_3), is solubilized in hot water in a jacketed vessel (3). The proportions of the algae and salt are, per 2 liters of water: 1 Kg of ground *Macrocystis* and 10 g of K_2CO_3 . The perforated basket (16) of freshly-ground *Macrocystis* is immersed in a jacketed vessel (3). The dry, ground algae *Ulva* is added and mixed. The solution pH is controlled (near to pH 10), at a temperature of $65 \pm 5^\circ\text{C}$, for a time of 2 hr, under constant stirring (mixer 17), for obtaining a high extraction yield of potassium alginate. In this step, a neutralization reaction occurs between the alginic acid comprised in the algal particles and the potassium alkali, which produces potassium alginate in aqueous solution.

[0015] The reaction carried out is as follows:



[0016] This reaction is important since to a large extent the yield and quality of the final product depends on the control of the physical-chemical parameters that are involved therein.

- **pH measurement:** Immediately after the digestion step is finished, the pH is reduced with an acid agent, phosphoric acid (H_3PO_4) from pH of about 9-5-10 to pH 4.5 - 6.2 at a temperature of about 50°C, for achieving better stabilisation of the final product.
- **Filtering:** Using a plate and frame filter-press (7) using a cellulose media as a filtering media. An objective of the filtration is to clarify the pH-adjusted extract solution by removing the non-soluble particles that remain after the digestion.
- **Storage:** Using fiberglass vats (8, 9, 10 and 11) the fertilizer solution product is allowed to cool for approximately between 1 to 2 days. Storage includes avoiding any kind of contamination, damage or deterioration of the product, while it is not dispatched, and controlling its environmental conditions: temperature and exposure to light.
- **Packing:** The liquid fertilizer complement product is packaged into plastic containers for its subsequent transport and distribution. It is necessary to note the importance of the packaging as an aspect of the quality in the product delivery, since it is the main way of maintaining quality in the distributed products.

[0017] The fertilizer complement can further comprise mineral elements, aminoacids, vitamins and auxin- and cytokinin-type phytohormones, based on the marine algae *Ulva* and *Macrocystis* contributions.

[0018] The fertilizer complement according to the present invention comprises algal particles as well as some inorganic compounds as indicated in table 1, wherein are showed the percentages of these components on 100 L of the final product.

Table 1

Ingredients	%
<i>Macrocystis</i> and <i>Ulva</i> algal particles	56
Water	43
hydrochloric acid	<0.01
potassium carbonate	0.6
phosphoric acid	0.3
potassium sorbate	0.1

Percentages calculated in basis of 100 L of the final product.

[0019] The obtained product, prepared from "organic" sources and the process above described, is able to provide plant growing certifiable for for being used in organically grown food crops. Thus the fertilizer complement can be added to plants growing media in order to provide a growth stimulator source which complies with "organic" standards for organically grown food.

[0020] As stated above, it is highly desirable to obtain a product which meets with the requirements established in the organic standards: EU N° 2092/1; USDA/NOP-Final rule (USA), and the JAS Japanese Agricultural standard for Organic Agricultural Products (Japan).

[0021] Thus, as an organic product for being applied to organically grown food, it is necessary to guarantee that the obtained final product is free of toxic elements, specifically heavy metals, which are hazardous for living beings. Table 2 shows the level of arsenic, mercury and cadmium contained in the final product according to the present invention, wherein said levels have been measured by atomic absorption spectroscopy.

Table 2

HEAVY METALS COMPOSITION FOR THE ORGANIC COMPLEMENT FERTILIZER			
	Arsenic*	Mercury*	Cadmium*
Maximum allowed by the EEC	(mg/kg)	(mg/kg)	(mg/kg)
<i>Ulva</i>	1.03	0.01	0.91
<i>Macrocystis</i>	0.24	0.00	0.24
Final product	0.05	0.00	0.02

* Atomic absorption spectroscopy, mean value of duplicated samples

[0022] According to the mentioned standards, the heavy metals content, specifically As, Hg, and Cd, must not be higher than 5%. The product of the instant invention fulfills satisfactorily these requirements.

Example:

[0023] In a preferred embodiment of the invention, a vessel is provided with water (100 L) at a temperature of about 40-60°C and 1.25 liters of 0.2 N HCl. 50 kg of minced *Macrocystis* is added and stirred during 30-40 minutes. The residual liquid is removed and the solid is ground and then incorporated into a second vessel with water (200 liters), potassium carbonate (1 kg) and minced *Ulva* (50 kg) for 2 to 2-1/2 hours at a temperature between 60-70°C. After filtering, the pH is lowered to a pH between 4.5 - 6.2 with phosphoric acid, to produce the resulting liquid fertilizer complement.

Claims

1. A process for making a fertilizer complement, wherein an organic vegetal growth stimulator is obtained from *Ulva* and *Macrocystis* algae, comprising the steps of acidifying the *Macrocystis* with an acidifying agent, and digesting the acidified *Macrocystis* and the *Ulva* with an alkalinizing agent.
2. The process according to claim 1, wherein the growth stimulator comprises auxin-type and cytokinin-type phyto-hormones.
3. The process according to claim 1 wherein the the *Ulva* algae comprises *Ulva rigida* and the *Macrocystis* comprises *Macrocystis pyrifera*.
4. The process according to claim 1, wherein the acidifying agent comprises 0.2 N hydrochloric acid.
5. The process according to claim 1, wherein the alkalinizing agent is potassium carbonate (K_2CO_3).
6. The process according to claim 1, comprising the steps of:
 - 1) receiving fresh algae *Macrocystis* and receiving algae *Ulva*,
 - 2) grinding the alga *Ulva*,
 - 3) weighing the fresh *Macrocystis* and *Ulva* algae,
 - 4) washing the fresh *Macrocystis* in water,
 - 5) mincing the washed *Macrocystis*,
 - 6) treating the minced *Macrocystis* with an acidifying agent,
 - 7) draining the residual liquid and separating the solid, minced *Macrocystis*,
 - 8) optionally washing the solid, minced *Macrocystis* with cold water,
 - 9) optionally grinding the washed *Macrocystis*,
 - 10) digesting the freshly ground *Macrocystis* with a potassium carbonate solution under stirring and admixing the dry ground *Ulva*,
 - 11) adjusting the pH of the digested mixture to acidic with phosphoric acid,
 - 12) filtering the digested mixture;
 - 13) optionally storing the filtered solution for 1 to 2 days, and
 - 14) optionally packaging the resulting product in plastic containers.
7. The process according to claim 6, wherein the weight ratio of the *Macrocystis* to *Ulva* is 1:1.
8. The process according to claim 7, wherein the weight ratio of *Macrocystis* to *Ulva* is up to 10:1 when the *Ulva* is in a dried state.
9. The process according to claim 7, wherein the step of washing results in a low electric conductivity in the washed *Macrocystis*.
10. The process according to claim 6, wherein the step of grinding the *Ulva* reduces the size to about 0.2 mm.
11. The process according to claim 6, wherein the particle size of the *Macrocystis* in step 5) is about 1 cm and in step

9) is about 0.4 cm.

12. The process according to claims 4, wherein the ratio are: water (200 L): *Macrocystis* (100 kg): 0.2 N HCl (2.5 l).

13. The process according to claim 12, wherein the acid treatment is carried out with 0.2N hydrochloric acid during 30-40 min at 50°C with constant stirring.

14. The process according to claims 5 and 6, **characterized in that** the ratio are: water (400 l): algae mixture (200 kg): K₂CO₃ (2 kg).

15. The process according to claim 14, **characterized in that** the alkaline treatment is carried out with K₂CO₃ during 2 hours at 65 ± 5 °C with constant stirring.

16. The process according to claim 15, **characterised in that** the digestion achieves a pH value of about 10 ± 0.2, allowing to obtain high extraction yields of potassium alginate.

17. The process according to claim 6, wherein the final pH value of the product must be regulated with technical degree phosphoric acid (H₃PO₄).

18. The process according to claim 14, **characterized in that** the final pH value of the product must be comprised in the range of about 4.5 and 6.2 in order to achieve a better stabilisation.

19. The process according to claim 6, **characterized in that** the final product must be filtered through a plate and frame press-filter (7) using as filtration media cellulose covers.

20. A fertilizer complement obtained by the process of claim 1.

21. The fertilizer complement according to claim 20, further comprising mineral elements, aminoacids, vitamins and auxin- and cytokinin-type phytohormones based on the marine algae *Ulva* and *Macrocystis* contributions.

FIGURE 1: FERTILIZATION ORGANIC COMPLEMENT FLOW-CHART






FIGURE 2: FLOW-CHART: MASS BALANCE FOR THE MANUFACTURE OF THE FERTILIZATION ORGANIC COMPLEMENT



EP 1 437 334 A1

FIGURE 3: THREE-DIMENSIONAL PHOTOGRAPHY OF THE EQUIPMENT FOR THE ORGANIC COMPLEMENT PRODUCTION.





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 04 29 0051

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	EP 0 855 375 A (ENGRAIS COMPOSES MINERAUX ET A) 29 July 1998 (1998-07-29) * page 1, line 28 - line 32 * * example 1 * * claims 1,2 *	1-21	C05F11/10
X	US 4 699 644 A (BRANDT ALAN E ET AL) 13 October 1987 (1987-10-13) * column 2, line 24 - line 38 *	20,21	
A		1-19	
X	EP 0 132 360 A (STAUFFER CHEMICAL CO) 30 January 1985 (1985-01-30) * page 4, line 21 - line 24 * * claims 3,4 *	20,21	
A		1-19	
X	DD 291 684 A (LPG PFLANZENPRODUKTION THOMAS) 11 July 1991 (1991-07-11) * page 2, line 4,5 *	20,21	
A		1-19	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C05F
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 8 April 2004	Examiner Saldamli, S
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 (03.02.92) (P06-C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 04 29 0051

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-04-2004

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0855375	A	29-07-1998	FR	2758816 A1	31-07-1998
			AT	205819 T	15-10-2001
			DE	69801688 D1	25-10-2001
			DE	69801688 T2	20-06-2002
			DK	855375 T3	21-01-2002
			EP	0855375 A2	29-07-1998
			ES	2165131 T3	01-03-2002
			PT	855375 T	28-02-2002

US 4699644	A	13-10-1987	NONE		

EP 0132360	A	30-01-1985	DK	352384 A	23-01-1985
			EP	0132360 A2	30-01-1985
			ES	534511 D0	01-10-1985
			ES	8600003 A1	01-01-1986
			JP	60054678 A	29-03-1985
			ZA	8405579 A	29-05-1985

DD 291684	A	11-07-1991	DD	291684 A5	11-07-1991

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)